Mon. Not. R. Astron. Soc. **000**, ??–?? ()   Printed 5 February 2008   (MN LaTeX style file v2.2)

arXiv:0802.0390v1 [astro-ph] 4 Feb 2008

Constraints on fall-back disks in radio pulsars and anomalous X-ray pulsars

P. B. Jones*
University of Oxford, Department of Physics, Denys Wilkinson building, Keble Road, Oxford OX1 3RH, England

ABSTRACT
Calculations have been made of fall-back disk heating by the pulsar wind as distinct from the soft X-rays emitted by the neutron-star surface. The relation between these heating rates and measured near-infrared fluxes in the K and K_s-bands places severe constraints on the inner radii of any fall-back disks that may be present in radio pulsars and in some anomalous X-ray pulsars. The lower limits found are so large that the disks concerned can have no significant effect on pulsar spin down.

Key words: accretion, accretion disks - stars: neutron - pulsars: general.

1 INTRODUCTION

Dust and debris disks are a commonplace phenomenon in astrophysics, being a feature of the major planets and of pre-main-sequence stars (see, for example, Beckwith et al 1990). The properties of dust disks or of viscous ionized disks formed as a consequence of fall-back in supernovae have been a matter of continuing interest in neutron-star physics. But until recently, there has been no observational evidence that they exist. For a bibliography, we refer to Perna, Hernquist & Narayan (2000), Menou, Perna & Hernquist (2001), Blackman & Perna (2004), Ekşi & Alpar (2005), Ekşi, Hernquist & Narayan (2005) and to the many earlier papers on the subject cited by these authors.

Further interest has been stimulated by flux measurements on the anomalous X-ray pulsar 4U 0142+61 in the 4.5 and 8.0 μm infrared bands (Wang, Chakrabarty & Kaplan 2006). The energy spectrum derived from these measurements, and from earlier K_s-band and optical fluxes, is consistent with blackbody emission from a dust disk with an inner radius surface temperature of 1200 K, superimposed on a power-law spectrum. The optical emission, which is pulsed at the neutron-star rotation frequency, consists almost entirely of the power-law component, which is assumed to be of magnetospheric origin.

There are also flux limits at 4.5, 8.0 and 24 μm in the mid-infrared for a limited number of isolated neutron stars (Wang, Kaspi & Higdon 2007) and for some supernova remnants at 4.5 and 8.0 μm (Wang, Kaplan & Chakrabarty 2007). But in the K and K_s-bands, a considerable number of magnitudes have been measured. These have been summarized by Mignani et al (2007) and include radio pulsars, anomalous X-ray pulsars (AXP), and soft-gamma repeaters (SGR). The purpose of the present paper is to see how these measurements constrain possible fall-back disk parameters.

Fall-back disk formation is a detail of the supernova event and no attempt is made to describe it here. Initially, the disk must have been internally ionized and hence viscous, with mass and angular momentum transfer rates intrinsic to its formation. We assume that at some later time, a transition to a passive disk occurs through the the thermal ionization instability (see Menou et al 2001) in which free electron recombination produces a sudden decrease in opacity which propagates rapidly inward. The disk formed has little internal ionization and viscosity, and is gaseous. Its luminosity is then principally a result of interaction with the pulsar wind or with blackbody radiation from the neutron star surface. The distribution of mass between dust grains and gaseous molecules depends on the various sublimation temperatures concerned. Our assumption is that, at the time of observation, the inner radius of the disk lies outside the light-cylinder radius R_{LC} of the neutron star. An important consequence is that we can treat the disk as a partial termination of the pulsar wind rather than as a boundary condition to be satisfied by the solution for the magnetospheric fields.

A previous paper (Jones 2007) on the interaction of a thin passive disk with the pulsar wind described the ablation processes in some detail and calculated the alignment and precession torques acting on the disk. Section 2 of the present paper extends the discussion of disk luminosity. The K and K_s-band magnitudes listed by Mignani et al for sets of radio pulsars and AXP or SGR are used, in Section 3, to constrain the inner radii and temperatures of passive fall-back disks that might be present in these objects. Where possible, the predicted mid-infrared fluxes from these objects are compared with the flux limits found by Wang, Kaspi & Higdon.

---

* E-mail: p.jones1@physics.ox.ac.uk

2 INTERACTION WITH THE PULSAR WIND

The Deutsch vacuum solution for the electromagnetic fields **E** and **B** of a rotating neutron star (see Michel & Li 1999, Ekşi & Alpar 2005) makes it possible to write down explicit expressions for the spherical polar components of the momentum density **p**,

$$\mathbf{p} = \frac{1}{4\pi c}\mathbf{D} \times \mathbf{B}, \tag{1}$$

in the inertial frame at radii $r > R_{LC}$, the z-axis being parallel with the neutron-star spin angular velocity $\mathbf{\Omega}$. The momentum density in a physical pulsar wind, in which both fields and relativistic particles are present, may differ from this in detail, but the time-averaged long-range part of the azimuthal component $\langle \mathbf{p}_\phi \rangle$ must be finite because it is directly related to the neutron-star spin-down torque (see Section 2.1 of Jones 2007). The composition of the wind does not significantly affect its interaction with the disk and it is therefore correct to use this quantity derived from the vacuum solution in calculations of ablation and, in particular, of alignment and precession torques.

The vacuum-solution time-averaged polar component is $\langle \mathbf{p}_\theta \rangle = 0$ for all angles, but its uni-directional time average, taken in the direction of either increasing or decreasing θ, can contribute to heating of the disk surface though not to a torque. With the addition of this component, the power input per unit area, to the one side of a thin disk that is viewed by an observer, can be written down and equated with the blackbody radiation emitted at the local temperature $T(r)$,

$$\sigma T^4(r) = \frac{3\mathcal{L}_w R_{LC}}{8\pi^2 r^3}\left(\sin\beta + \frac{2}{3}\cot\xi\right), \tag{2}$$

where $\mathcal{L}_w$ is the pulsar wind luminosity, β is the disk tilt angle and ξ is the angle between the neutron-star magnetic and spin axes. The first term in equation (2) is derived from the long-range part of $\langle \mathbf{p}_\phi \rangle$ by means of equation (14) of Jones (2007). The second term, in the approximation of small β, is from the long-range part of the uni-directional time average of $\mathbf{p}_\theta$ evaluated in the vicinity of $\theta = \pi/2$. The status of the vacuum solution form of this term is much less certain. Apart from an apparent unphysical singularity at $\xi = 0$, which is a consequence of the vacuum field solution and our use of $\mathcal{L}_w \propto \sin^2\xi$ as a working parameter, the term is linearly dependent on the radial component of the electric displacement **D** which is undetermined because the total charge of the star and the physical, as opposed to vacuum, magnetosphere is an unknown quantity, as has been emphasized by Michel & Li (1999).

The values that might be found for the tilt angle β are, of course, unknown, but there is a case for assuming that its value at formation is non-zero as a consequence of the fact that type II supernovae are generally asymmetric (see Goldreich, Lai & Sahrling 1997). This suggests that the angular momenta of the neutron star, the fall-back disk and the ejected matter may not all be precisely parallel.

The uncertainties in β and in the polar term are so considerable that we shall replace the vacuum-field quantity $\sin\beta + 2/3\cot\xi$ in equation (2) by a free parameter ζ and adopt a conservative range of values $0.03 \leqslant \zeta \leqslant 0.3$. Neglect of the wind albedo is a further uncertainty in equation (2), but not one that is serious because relativistic particles either in the wind or accelerated in the diffuse surface of the disk as described previously (Jones 2007) are attenuated only at depths of $10^2 - 10^3$ g cm^{-2} within the disk. (We assume that the mean disk density $\bar{\Sigma}$ exceeds 10^3 g cm^{-2}.) For the same reason, it is not anticipated that infrared or optical emission from an undistorted disk would have an observable component pulsed at the neutron-star rotation period.

We anticipate that, even for young neutron stars, almost all of the potential disk ablation will have occurred by the time of observation. Interaction with the pulsar wind increases the inner radius r_i which is given by

$$r_i^3(t) = \frac{3cIm_H\mathcal{N}_\epsilon\zeta(\Omega_0 - \Omega(t))}{4\pi^2\bar{\Sigma}}, \tag{3}$$

which is valid provided the solution $r_i(t)$ exceeds the real r_i at formation. Here Ω_0 is the neutron-star spin angular velocity at formation, I is the neutron-star moment of inertia and m_H the mass of the hydrogen atom. The remaining parameter $\mathcal{N}_\epsilon$ is the number of baryons (mostly neutrons) removed from the disk per unit incident wind energy (see Jones 2007; equations 17 and 18). The mean mass density $\bar{\Sigma}$ is the initial value at formation. Owing to the number of unknown parameters it contains, equation (3) does not provide a useful constraint on disk inner radii derived from equation (2). But it shows that, in most cases, the existing r_i must be near its $\Omega = 0$ limit. Because equation (3) is based on the interaction of the disk with the pulsar wind, it ceases to be valid when the light-cylinder radius exceeds r_i. But for the radio pulsars, this occurs only for Ω so small as to be of no interest.

3 DISK PARAMETERS

Recent observations in the near-infrared have produced H, K and K$_s$-band magnitudes for many isolated neutron stars, including both radio pulsars and AXP. A useful listing of these has been published by Mignani et al (2007) who also include estimates of the distances d. We have used the Spitzer Science Centre magnitude to flux-density converter to obtain the observed energy fluxes per unit frequency interval f_ν. Equation (2) provides the disk surface temperature for each object as a function of radius and it is then straightforward, by integration and interpolation and with a number of moderate assumptions, to find the disk inner radius that gives f_ν. In particular, we assume that the angle $\theta_\perp$ between the disk normal and the line of sight is fixed and given by $\cos\theta_\perp = 0.5$. No attempt has been made to correct for the small K-band interstellar extinction. The values found for r_i are quite insensitive to the outer radius r_o because, for the temperatures found here, the outer part of the disk contributes little to the K-band flux. Therefore, we have adopted a fixed $r_o = 3.4r_i$, equal to the best-fit ratio found by Wang et al in the case of 4U 0142+61. With these assumptions, values of r_i and the inner radius temperature T_i have been computed and are given in the right-hand six columns of Table 1 for $\zeta = 0.03, 0.1, 0.3$.

The disks of Table 1 need not be massive: for $\zeta = 0.03$, the masses are in the interval 10^{-7} - 10^{-5} M$_\odot$, assuming a density $\bar{\Sigma} = 10^3$ g cm^{-2}, but their angular momenta can be of the same order as that of the neutron star. Certainly for the lower temperatures given in Table 1, the disks must consist almost entirely of dust grains and only a small gaseous component. But even a very low concentration of free electrons and ions is sufficient to stop the pulsar wind (see Jones 2007) and there can be no doubt that such concentrations must be present from a variety of sources.

For consistency with our Section 1 definition of a fall-back disk, we require that a satisfactory solution has $r_i > R_{LC}$. The first six rows in the Table refer to radio pulsars, and they satisfy this condition by large margins. But it is interesting to see that, in many cases, the AXP and SGR do not. In particular, there is no solution for 4U 0142+61, the object for which evidence of a disk is most compelling. This is not a cause for concern because equation (2) contains only the pulsar wind contribution to disk heating. For the AXP and SGR, the soft X-ray luminosity $\mathcal{L}_{bol}$, possibly of thermal origin, exceeds $\mathcal{L}_w$ by several orders of magnitude so that its power input can be the more important if the disk is not of negligible thickness but subtends a finite solid angle at the neutron star, as in the model adopted by Wang et al (2006).

It is reasonable to assume that a disk, immediately after the ionization instability transition, would be plane and thin. If there is a gaseous component, it would have a diffuse surface of approximate depth

$$h = \sqrt{\frac{k_B T}{m_{A,Z}\Omega_K^2}}, \qquad (4)$$

where $m_{A,Z}$ is an atomic (or molecular) mass and Ω_K is the Kepler angular frequency. For the 4U 0142+61 best-fit inner radius $r_i = 2.0 \times 10^{11}$ cm found by Wang et al, the very small Kepler frequency, 1.5×10^{-4} rad s^{-1} leads to a fairly large depth, $h \approx 2 \times 10^9$ cm, for atomic hydrogen at $T = 10^3$ K. The 4U 0142+61 luminosity is $\mathcal{L}_{bol} \sim 10^{35}$–$10^{36}$, so that although the disk is thin, the X-ray power input, $(1-\eta_d)h\mathcal{L}_{bol}/r_i$ exceeds that given by equation (2) even for the large value of the X-ray albedo η_d adopted by Wang et al (2006). The solid angle subtended by the disk may also be increased by departures from the plane assumption arising from r-dependent alignment torques (Jones 2007) or disk instability (Petterson 1977; Pringle 1996). The X-ray contribution to disk heating is therefore important for the AXP and SGR, but not for the radio pulsars which have $\mathcal{L}_{bol} \ll \mathcal{L}_w$.

4 COMPARISON WITH OBSERVATION

The electromagnetic spectra of a limited number of radio pulsars are known over wide frequency intervals. In the case of B0656+14, as an example, apart from the lack of measurements between 10^{10} and 10^{14} Hz, the shape of the spectrum is of blackbody emission from the neutron-star surface superimposed on a background which decreases monotonically with frequency and is possibly of some common magnetospheric origin (see Koptsevich et al 2001; Fig. 6) The K-band flux listed in Table 1 is consistent with this spectrum. It is also known that the optical emission is pulsed at the rotation frequency. For example, Shearer et al (1997), observing B-band emission, found a 1σ upper limit of only 8×10^{-31} erg s^{-1} cm^{-2} Hz^{-1} on the unpulsed component of the flux. With regard to the AXP listed in Table 1, the optical emission from 4U 0142+61 has a pulsed fraction of 0.3, and the optical and X-ray pulse profiles are in phase and of similar shape (Kern & Martin 2002, Dhillon et al 2005).

Unambiguous evidence for the presence of a disk would be an unpulsed blackbody excess which, for the temperatures given in Table 1, would probably be best observed in the 4.5 and 8.0 μm bands. With the possible exception of 4U 0142+61, existing spectra provide no such evidence.

Even for the smallest value of ζ assumed in Table 1, the radio pulsar disk inner radii that would be compatible with the K-band fluxes are all larger than the best-fit $r_i = 2.0 \times 10^{11}$ cm found by Wang et al (2006) in the case of 4U 0142+61, although they are of a similar order of magnitude. It might be argued that the correct value of ζ is much smaller than 0.03, perhaps by one or more orders of magnitude. For this to be so, the neutron star spin $\mathbf{\Omega}$ would have to be almost exactly normal to the plane of the disk. Also, the star and magnetosphere must be electrically neutral to the extent that the uni-directional time-average of $\mathbf{p}_\theta$, evaluated at $\theta = \pi/2$, is two or more orders of magnitude smaller than the vacuum solution value. Instead, we suggest that the inner radii given for the radio pulsars in Table 1 are broadly correct as lower limits.

Heating of a disk with inner radius $r_i > R_{LC}$ by the pulsar wind is capable of producing the observed K_s-band fluxes in only two of the AXP listed in Table 1. But these objects are differ significantly from radio pulsars in having very high soft X-ray luminosities, $\mathcal{L}_{bol} \gg \mathcal{L}_w$, which give most of the disk heating, as in the model adopted by Wang et al (2006). The two exceptional AXP, J1048-5937 and J1810-197, are interesting in that 3σ upper limits for their mid-infrared fluxes at 4.5, 8.0 and 24 μm have been published recently by Wang, Kaspi & Higdon (2007). These fluxes can also be calculated from the r_i and T_i values given in the Table. For J1810-197, all the predicted fluxes are below the the 3σ upper limits. However, the J1048-5937 4.5 μm 3σ upper limit of 8×10^{-29} erg cm^{-2} s^{-1} Hz^{-1} is exceeded by the predicted flux of 1.2×10^{-28} erg cm^{-2} s^{-1} Hz^{-1} for $\zeta = 0.03$ which increases to 3.1×10^{-28} erg cm^{-2} s^{-1} Hz^{-1} at $\zeta = 0.3$. A very small value, $\zeta = 0.012$ is required to fit both the measured K$_\mathrm{s}$-band flux and the published 3σ upper limit. In this case, the disk is at a higher temperature $T_i = 1300$ K and has a smaller radius $r_i = 7.9 \times 10^{10}$ cm. But an alternative possibility is that the source of both the K$_\mathrm{s}$-band and 4.5 μm fluxes is magnetospheric rather than a heated disk, which could be confirmed by a search for a pulsed component.

5 CONCLUSIONS

The present paper considers heating of a thin passive disk by non-radial components of the pulsar wind and computes inner radii and temperatures that are compatible with the observed K and K$_\mathrm{s}$-band fluxes for a number of radio pulsars and AXP. In the case of the radio pulsars, the inner radii are two or more orders of magnitude larger than the R_{LC} and the disks concerned, if they exist, can have no significant effect on pulsar spin-down. The conservative, and probably incorrect, assumption made is that the observed K and K$_\mathrm{s}$ fluxes have no pulsed component and are of thermal origin. It is also assumed that there is no internal disk viscosity and that heating by the radial flux of soft X-rays from the

Table 1. Disk inner radii r_i that are consistent with measured K and K_s-band magnitudes have been calculated, under the assumptions made in Section 3, for a number of radio pulsars and AXP or SGR. Inner radii temperatures T_i are also given. For J0142+61, J1810-197, IE 2259+586, and SGR 1806-20, the values of the wind luminosity $\mathcal{L}_w$ have been obtained from the periods P and spin-down rates given, respectively, by Gavril & Kaspi (2002), Ibrahim et al (2004), Woods et al (2004) and Mereghetti et al (2005). For the remaining neutron stars, the periods and $\mathcal{L}_w$ are from the Australia Telescope National Facility Pulsar Catalogue (Manchester et al 2005). The distances d are those listed by Mignani et al (2007). The energy flux densities f_ν have been obtained from the observed K or K_s-band magnitudes using the Spitzer Science Centre magnitude to flux-density converter. The right-hand six columns give r_i and T_i for $\zeta = 0.03, 0.1, 0.3$. Blank spaces indicate the absence of a solution with $r_i > R_{LC}$.

neutron star	P s	$\mathcal{L}_w$ erg s^{-1}	f_ν erg cm^{-2} s^{-1} Hz^{-1}	d kpc	r_i cm $\zeta = 0.03$	T_i K	r_i cm $\zeta = 0.1$	T_i K	r_i cm $\zeta = 0.3$	T_i K
B0531+21	0.033	4.6×10^{38}	2.0×10^{-26}	1.73	4.6×10^{11}	1970	1.0×10^{12}	1440	2.0×10^{12}	1180
B0633+17	0.237	3.2×10^{34}	2.9×10^{-30}	0.16	2.5×10^{11}	470	4.0×10^{11}	440	6.2×10^{11}	420
B0656+14	0.385	3.8×10^{34}	6.1×10^{-30}	0.29	2.6×10^{11}	540	4.2×10^{11}	500	6.6×10^{11}	470
B0833-45	0.089	6.9×10^{36}	2.0×10^{-29}	0.29	1.0×10^{12}	480	1.7×10^{12}	450	2.6×10^{12}	420
B1509-58	0.151	1.8×10^{37}	1.2×10^{-28}	4.2	6.2×10^{11}	1020	1.1×10^{12}	880	1.9×10^{12}	780
J1119-6127	0.408	2.3×10^{36}	$< 1.1 \times 10^{-29}$	6.0	$> 5.5 \times 10^{11}$	< 860	$> 9.7 \times 10^{11}$	< 760	$> 1.6 \times 10^{12}$	< 690
J0142+61	8.69	1.2×10^{32}	6.1×10^{-29}	$\geqslant 5$	-	-	-	-	-	-
J1048-5937	6.45	5.6×10^{33}	2.0×10^{-29}	3	1.3×10^{11}	1110	2.4×10^{11}	950	4.1×10^{11}	840
J1708-4009	11.00	5.8×10^{32}	3.2×10^{-28}	5	-	-	-	-	-	-
J1810-197	5.54	2.7×10^{33}	3.2×10^{-29}	4	5.0×10^{10}	1840	1.1×10^{11}	1380	2.0×10^{11}	1140
IE 2259+586	6.98	5.6×10^{31}	1.4×10^{-29}	3.0	-	-	-	-	4.2×10^{10}	1510
SGR 1806-20	7.56	5.0×10^{34}	6.1×10^{-29}	15	-	-	1.7×10^{11}	2180	4.0×10^{11}	1570

neutron-star surface is negligible except in the case of the AXP.

The possibility that some of the individual neutron stars listed in Table 1 have disks has been investigated by previous authors. K-band fluxes from disks with inner radii near the Alfvén surface have been calculated in the case of B0656+14, and the four AXP J0142+61, J1048-5937, J1708-4009 and IE 2259+586, by Perna, Hernquist & Narayan (2000). Also, the radio pulsars B0531+21 and B0833-45 were considered by Blackman & Perna (2004). The disk model differs from that of Section 2 in that the sources of heating were restricted to internal viscous dissipation and the soft X-ray spectrum of the neutron-star surface. Even so, the predicted K-band fluxes for the four AXP were $f_\nu \sim 10^{-27} - 10^{-26}$ erg cm^{-2} s^{-1} Hz^{-1}, considerably higher than the observed values listed in Table 1. In the case of the three radio pulsars, inclusion of heating by non-radial wind components is crucial and leads us to the conclusion that the proposed disks are not consistent with the observed K-band fluxes. This is also true in the general case, treated by Ekşi & Alpar (2005), of disks with inner radii near the light-cylinder radius. This conclusion should perhaps be qualified because we do not consider disks with $r_i < R_{LC}$. Thus a disk satisfying this condition at formation, with R_{LC} given by the initial angular frequency Ω_0, would have its middle section removed by wind ablation leaving an outer section with the properties given in Table 1 and an inner section completely within the present light-cylinder radius. This inner sector would form part of the closed magnetosphere and would not intersect open magnetic flux lines or interact with the pulsar wind. The properties of such remnant disks, if they exist, are not considered in this paper.

REFERENCES

Beckwith S. V. W., Sargent A. I., Chini R. S., Güsten R., 1990, AJ, 99, 924

Blackman E. G., Perna R., 2004, ApJ, 601, L71

Dhillon V. S., Marsh T. R., Hulleman F., van Kerkwijk M. H., Shearer A., Littlefair S. P., Gavril F. P., Kaspi V. M., 2005, MNRAS, 363, 609

Ekşi K. Y., Alpar M. A., 2005, ApJ, 620, 390

Ekşi K. Y., Hernquist L., Narayan R., 2005, ApJ, 623, L41

Gavril F. P., Kaspi V. M., 2002, ApJ, 567, 1067

Goldreich P., Lai D., Sahrling M., 1997, Unsolved Problems in Astrophysics, ed. Bahcall J. N., Ostriker J. P., Princeton University Press, Princeton, N. J., p. 269

Ibrahim A. I. et al, 2004, ApJ, 609, L21

Jones P. B., 2007, MNRAS, 382, 871

Kern B., Martin C., 2002, Nat., 417, 527

Koptsevich A. B., Pavlov G. G., Zharikov S. V., Sokolov V. V., Shibanov Yu. A., Kurt V. G., 2001, A&A, 370, 1004

Manchester R. N., Hobbs G. B., Teoh A., Hobbs M., 2005, AJ, 129, 1993

Menou K., Perna R., Hernquist L., 2001, ApJ, 559, 1032

Mereghetti S. et al, 2005, ApJ, 628, 938

Michel F. C., Li H., 1999, Phys. Rep., 318, 227

Mignani R. P., Perna R., Rea N., Israel G. L., Mereghetti S., Lo Curto G., 2007, A&A, 471, 265

Perna R., Hernquist L., Narayan R., 2000, ApJ, 541, 344

Petterson J. A., 1977, ApJ, 216, 827

Pringle J. E., 1996, MNRAS, 281, 357

Shearer A. et al, 1997, ApJ, 487, L181

Wang Z., Chakrabarty D., Kaplan D. L., 2006, Nat., 440, 772

Wang Z., Kaplan D. L., Chakrabarty D., 2007, ApJ, 655, 261

Wang Z., Kaspi V. M., Higdon S. J. U., 2007, ApJ, 665,

1292
Woods P. M. et al, 2004, ApJ, 605, 378

This paper has been typeset from a TEX/ LATEX file prepared by the author.